



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Also solved by H. A. LEVY, A. M. HARDING, J. A. COLSON, LEVI S. SHIVELY, H. C. FEEMSTER, ELMER SCHUYLER, and the PROPOSER.

391. Proposed by C. N. SCHMALL, New York, N. Y.

Show that the roots of the quadratic

$$ax^2 + 2bx + c = 0$$

are imaginary if a, b, c are in harmonic progression and have the same sign.

SOLUTION BY S. W. REAVES, University of Oklahoma.

Since a, b, c are in harmonic progression,

$$b = \frac{2ac}{a+c}.$$

The discriminant of the given quadratic then becomes

$$b^2 - ac = \left(\frac{2ac}{a+c} \right)^2 - ac = -ac \left(\frac{a-c}{a+c} \right)^2,$$

which is negative since a and c have the same sign. The discriminant being negative, the roots are imaginary.

Also solved by WALTER C. EELLS, ELMER SCHUYLER, F. M. MORGAN, H. C. FEEMSTER, J. A. COLSON, and A. M. HARDING.

Additional solutions of 387 were received from R. M. MATHEWS, HORACE OLSON, and G. Y. SOSNOW, after the December issue had gone to press.

Note. We have solutions of all problems proposed during 1913 in this section up to the September number, except 385, which was published in February.


GEOMETRY.

419. Proposed by S. LEFSCHETZ, University of Nebraska.

Given a circle and a tangent to it. To find a point on its circumference such that the sum of its distances to the tangent and its point of contact shall be equal to a given length.

SOLUTION BY S. W. REAVES, University of Oklahoma.

Let ORN be the given circle, OT the given tangent, and O the point of contact. Denote the given length by l .



Construction. Draw the diameter ON and extend it beyond N to P making $NP = 2l$. On OP as diameter construct the semi-circle $OSS'P$. On the given tangent lay off $OT = l$. Through T draw the line TSS' parallel to OP to meet the semi-circumference first in S . Drop a perpendicular from S to OP meeting OP in M and meeting the given circle in R and R' . Then R (or R') is the required point.

Proof. Let $QR = OM = x$, $OR = y$, $OS = z$, $ON = 2r$. Then by construction $OP = 2(r + l)$ and $MS = l$.

By elementary geometry $OM : OR = OR : ON$.

Hence

$$y^2 = 2rx. \quad (1)$$

Similarly

$$OM : OS = OS : OP.$$

Hence,

$$z^2 = 2x(r + l), \quad (2)$$

or, from (1),

$$z^2 = y^2 + 2lx. \quad (3)$$

But $z^2 = x^2 + l^2$, and therefore (3) becomes

$$y = \pm (l - x). \quad (4)$$

Since $y > 0$ and $l > x$, the lower sign is inadmissible. Hence $x + y = l$, or

$$QR + OR = l.$$

A solution exists when and only when $l \leq 4r$, and there is not more than one solution on the same side of the diameter through the point of contact.

Solutions were also received from WALTER C. EELLS, H. C. FEEMSTER, C. N. SCHMALL, ELMER SCHUYLER, and A. H. HOLMES.

420. Proposed by C. N. SCHMALL, New York City.

Four spheres are described so that each touches a face of a given triangular pyramid and the other three faces produced. If the radii of the escribed spheres be r_1, r_2, r_3, r_4 , and r be the radius of the inscribed sphere, show that

$$\frac{1}{r_1} + \frac{1}{r_2} + \frac{1}{r_3} + \frac{1}{r_4} = \frac{2}{r}.$$

SOLUTION BY HORACE OLSON, Chicago, Illinois.

Let $ABCD$ be a given tetrahedron.

Let the faces ABC, ABD, ACD , and BCD be represented by small letters corresponding to the opposite vertices of the tetrahedron, *i. e.*, by d, c, b , and a respectively.

Let V represent the volume of the tetrahedron $ABCD$.

Let O be the center, and r the radius, of the inscribed sphere.

Let O_1, O_2, O_3 , and O_4 be the centers, and r_1, r_2, r_3 , and r_4 the radii, of the escribed spheres opposite to A, B, C , and D , respectively.

Connect points O and O_3 with the 4 vertices of the tetrahedron.

Then,

$$V = AOBC + OBCD + OACD + OABD = \frac{ar}{3} + \frac{br}{3} + \frac{cr}{3} + \frac{dr}{3} = \frac{r}{3}(a + b + c + d),$$